



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/536, A61P 3/04, C07D 265/24, 498/04, 413/12 // (C07D 498/04, 265:00, 221:00)		A1	(11) International Publication Number: WO 00/40247
			(43) International Publication Date: 13 July 2000 (13.07.00)
(21) International Application Number: PCT/GB00/00032		John [GB/GB]; Alizyme Therapeutics Limited, 280 Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). (74) Agents: FLORENCE, Julia, Anne et al.; Kilburn & Strode, 20 Red Lion Street, London WC1R 4PJ (GB). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 6 January 2000 (06.01.00)			
(30) Priority Data: 9900413.7 8 January 1999 (08.01.99) GB 9917294.2 22 July 1999 (22.07.99) GB			
(71) Applicant (for all designated States except US): ALIZYME THERAPEUTICS LIMITED [GB/GB]; 280 Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB).			
(72) Inventors; and (75) Inventors/Applicants (for US only): HODSON, Harold, Francis [GB/GB]; 69 Whitcroft Way, Beckenham, Kent BR3 3AW (GB). DOWNHAM, Robert [GB/GB]; Cambridge Discovery Chemistry, The Merrifield Centre, Rosemary Lane, Cambridge CB1 3LQ (GB). MITCHELL, Timothy, John [GB/GB]; Cambridge Discovery Chemistry, The Merrifield Centre, Rosemary Lane, Cambridge CB1 3LQ (GB). CARR, Beverley, Jane [GB/GB]; Boys Bridge Cottage, Fen Road, Abbington Piggotts, Cambs. SG8 5PQ (GB). DUNK, Christopher, Robert [GB/GB]; Alizyme Therapeutics Limited, 280 Cambridge Science Park, Milton Road, Cambridge CB4 4WE (GB). PALMER, Richard, Michael,			


Published

With international search report.

(54) Title: 2-AMINO-4H-3,1-BENZOXAZIN-4-ONE DERIVATIVES FOR THE TREATMENT OF OBESITY

(57) Abstract

The use of a compound comprising formula (I) or a salt, ester, amide or prodrug thereof in the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality, e.g. in the control and inhibition of unwanted enzymes in products and processes. The compounds are also useful in medicine e.g. in the treatment of obesity and related conditions. The invention also relates to novel compounds within formula (I), to processes for preparing them and pharmaceutical compositions containing them. In formula (I) A is a 6-membered aromatic or heteroaromatic ring; and R¹ is a branched or unbranched alkyl (optionally interrupted by one or more oxygen atoms), alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, reduced arylalkyl, arylalkenyl, heteroaryl, heteroarylalkyl, heteroarylalkenyl, reduced aryl, reduced heteroaryl, reduced heteroarylalkyl or a substituted derivative of any of the foregoing groups.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LR	Sri Lanka	SE	Sweden		
EE	Estonia		Liberia	SG	Singapore		

2-AMINO-4H-3,1-BENZOXAZIN-4-ONE DERIVATIVES FOR THE TREATMENT OF OBESITY


The present invention provides known and novel compounds, their use in the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality (*in vivo*, as the enzyme naturally occurs), their use in medicine, and particularly in the prevention and/or treatment of obesity or an obesity-related disorder. Also provided are methods for the prevention and/or treatment of obesity or an obesity-related disorder and for promoting/ aiding non-medical weight loss and the use of the compounds in the manufacture of a medicament for the aforementioned indications. In respect of novel compounds the invention also provides processes for their manufacture, compositions containing them, and methods for manufacturing such compositions.

In the last 20 years, there has been an increasing trend in obesity in the populations of the developed world. The increased incidence of obesity is due in part to the ready availability of food in numerous retail outlets and westernised diets that have high saturated fat and lower fibre contents such that the food is energy dense. The lifestyle of the populations of the developed world has also become more sedentary with the increased mechanisation of society and the steady reduction of manual labour intensive industries. There now exists an energy imbalance between the energy intake from calorie dense foods and the reduced energy expenditure required for a sedentary lifestyle. Some of the excess energy intake is stored as fat in the adipose tissue, the accumulation of which over a period of time results in obesity and can be a significant contributory factor to other diseases and disorders.


Obesity is now recognised by the medical profession as a metabolic disease. In the USA, it is estimated that 25% of the adult population is considered clinically obese (Body Mass Index>30). Obesity can be a debilitating condition which reduces the quality of life and increases the risk of related disorders such as diabetes, cardiovascular disease and hypertension. It has been estimated that \$45 billion of US

healthcare costs, or 8% per annum of total healthcare spend, is as a direct result of obesity. The traditional approaches to long term weight management such as diet and exercise have proved ineffective alone to control the spread of obesity. Today, more than ever, there is considerable interest in developing safe, effective drugs for the treatment of obesity.

Pharmacological approaches to the treatment of obesity have focused on either developing drugs that increase energy expenditure or drugs that reduce energy intake. One approach to the reduction of energy intake is to reduce the body's ability to digest and absorb food, in particular fat. The key enzymes involved in the digestion of fat are hydrolytic enzymes. The most significant of the fat degrading enzymes are lipases, primarily, but not exclusively pancreatic lipase that is secreted by the pancreas into the gut lumen. The lipase inhibitor lipstatin has formed the basis of the anti-obesity drug, orlistat. Orlistat is the subject of published European Patent Application No. EP129748, which relates to compounds of formula:




where A is $-(CH_2)_5-$ or:



and their use in inhibiting pancreatic lipase and treating hyperlipaemia and obesity.

5 Orlistat has as its major active moiety a beta-lactone group that reacts to form an ester with the side chain hydroxyl group of serine 152 within the active site of pancreatic lipase. Even if orlistat provides an effective method for treating obesity, there remains a need to provide alternative drugs and methods for use in the control and treatment of obesity, obesity-related disorders and non-medical weight loss. Inhibitors of enzymes
10 involved in the degradation of fat are provided here and shown to be effective in the prevention and/or treatment of obesity, obesity-related disease and/or cosmetic weight loss.

US Patent No. 4,657,893 (Syntex) describes a broad class of 2-amino-4H-3,1-
15 benzoxazin-4-ones of the formula:




wherein R¹ is hydrogen or lower alkyl, R² and R³ are each independently hydrogen, halo, lower alkyl, hydroxy, lower alkoxy, lower thioalkyl, -NO₂, -N(R¹)₂, -NR¹COR¹,
25 -NHCON(R¹)₂ or NHCOOR¹; and X is *inter alia* -NHR where R is lower alkyl, lower alkenyl, lower alkynyl, optionally substituted lower cycloalkyl or optionally substituted phenyl lower alkyl. The compounds are said to be useful as serine protease inhibitors and to treat physiologic conditions and disease states known to involve serine proteases, or as contraceptives. The specification describes various conditions
30 and diseases involving enzymatic pathways, including inflammation, arthritis, tumor cell metastasis, pulmonary emphysema, mucocutaneous lymph node syndrome, adult

respiratory distress syndrome and pancreatitis. It is also suggested that the compounds may have antiparasitic, anticoagulant and/or antiviral activity. Similar compounds are also described by Krantz *et al* in *J. Med. Chem.* 1990 **33**:464-479.

5 2-Amino-4*H*-3,1-benzoxazin-4-ones as inhibitors of serine protease are also described by Hays *et al* in *J. Med. Chem.* 1998 **41**:1060-1067. This paper describes *inter alia* 2-(substituted phenyl)amino benzoxazinones, where the phenyl substituents include halogen, methyl, SMe, and OCF₃, as well as certain 2-(heterocyclic)amino benzoxazinones. Some of these compounds are also described in US Patent No.
10 5,652,237 (Warner Lambert).

German OLS 2315303 (Bayer AG) describes the preparation of compounds of the formula




where R is an alkyl or aryl residue which may be substituted by nitro, halogen, alkyl, alkoxy or an aryl group, and R' and R'' are each independently hydrogen, halogen, nitro, optionally substituted alkyl, cycloalkyl, aralkyl, aryl, alkoxy or aryloxy groups. The only values of R exemplified are nitrophenyl and mono- and di-chlorophenyl. The compounds are said to be useful as intermediates for pharmaceuticals and plant protection agents.

25

We have now found that a particular class of benzoxazinone compounds has activity as lipase inhibitors

Accordingly, a first aspect of the invention provides a compound comprising formula
30 (I)



(I)

5 or a pharmaceutically acceptable salt, ester, amide or prodrug thereof; in the manufacture of a medicament for the treatment of conditions which require the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality; wherein in formula (I):

10 A is a 6 membered aromatic or hetero-aromatic ring;

R¹ is a branched or unbranched alkyl (optionally interrupted by one or more oxygen atoms), alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, reduced arylalkyl, arylalkenyl, heteroaryl, heteroarylalkyl, heteroarylalkenyl, reduced aryl, reduced
 15 heteroaryl, reduced heteroarylalkyl or a substituted derivative thereof wherein the substituents are one or more independently chosen from the group consisting of halogen, alkyl, halosubstituted alkyl, aryl, arylalkyl, heteroaryl, reduced heteroaryl, reduced heteroarylalkyl, arylalkoxy, cyano, nitro, -C(O)R⁴, -CO₂R⁴, -SOR⁴, -SO₂R⁴, -NR⁶R⁷, -OR⁶, -SR⁶, -C(O)CX¹X²NR⁶R⁷, -C(O)NR⁴R⁵, -C(O)N(OR⁵)R⁶, -NR⁶C(O)R⁴,
 20 -CR⁶(NH₂)CO₂R⁶, -NHCX¹X²CO₂R⁶, -N(OH)C(O)NR⁶R⁷, -N(OH)C(O)R⁴, -NHC(O)NR⁶R⁷, -C(O)NHN⁶R⁷, -C(O)N(OR⁵)R⁶, or a lipid or steroid (natural or synthetic), with the proviso that any hetero atom substituent in R¹ and/or R² must be separated from the exocyclic nitrogen atom by at least two carbon atoms (preferably saturated); and

25

R^2 is hydrogen or is a group as defined above for R^1 ;

and where:-

5 R^4 is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, reduced heteroaryl, reduced heteroarylalkyl, $-OR^6$, $-NHCX^1X^2CO_2R^6$ or $-NR^6R^7$;

10 R^5 is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, reduced heteroaryl or reduced heteroarylalkyl;

R^6 and R^7 are independently selected from hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, reduced heteroaryl, heteroarylalkyl, reduced heteroarylalkyl or $-(CH_2)_n(OR^5)_m$ wherein n is 1 to 12, preferably 2 to 10, wherein m is 1-3 and R^5 is most preferably C_{2-10} alkyl; and

15 X^1 and X^2 are independently hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, reduced heteroaryl or reduced heteroarylalkyl.

20

In compounds of formula (I) any alkyl, alkenyl and alkynyl groups and moieties may be straight chain (unbranched) or branched chain. Straight chain alkyl, alkenyl and alkynyl groups or moieties may contain from 1 to 30 carbon atoms, eg. 1 to 25 carbon atoms, preferably 1 to 20 carbon atoms. Branched chain alkyl, alkenyl and alkynyl groups or moieties may contain from 1 to 50 carbon atoms, preferably 1 to 30 carbon atoms.

25

Preferred values for R^1 , R^4 , R^5 , R^6 , R^7 , X^1 and X^2 are as defined below for formulae (II) and (IIa). In particular, preferred values for R^4 , R^5 and R^6 are as defined for R^{13} and preferred values of R^7 are as defined for R^{14} hereinbelow.

30

In this text, 'reduced', in the context of 'reduced heteroaryl' and the like means fully or partially saturated.

5 Aryl groups include for example optionally substituted unsaturated monocyclic or bicyclic rings of up to 12 carbon atoms, such as phenyl and naphthyl, and partially saturated bicyclic rings such as tetrahydro-naphthyl. Examples of substituents which may be present on an aryl group include one or more of halogen, amino, nitro, alkyl, haloalkyl, alkoxy, phenoxy and phenoxy substituted by one or more of halo, alkyl or
10 alkoxy.


A heteroaryl group or moiety may be for example an optionally substituted 5- or 6-membered heterocyclic aromatic ring which may contain from 1 to 4 heteroatoms selected from O, N and S. The heterocyclic ring may optionally be fused to a phenyl
15 ring. Examples of heteroaryl groups thus include furyl, thienyl, pyrrolyl, oxazolyl, oxazinyl, thiazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, pyridyl, triazolyl, triazinyl, pyridazyl, pyrimidinyl, pyrazolyl, indolyl, indazolyl, benzofuranyl, benzothienyl, benzimidazolyl, benzoxazolyl, benzoxazinyl, quinoxaliny, quinoliny, quinazolinyl, cinnoliny, benzothiazolyl, pyridopyrrolyl. Suitable substituents include one or more
20 of halogen, oxo, amino, nitro, alkyl, haloalkyl, alkoxy, phenoxy and phenoxy substituted by one or more of halo, haloalkyl, alkyl or alkoxy.

A reduced heteroaryl group or moiety may be for example a fully or partially saturated derivative of the aforementioned heteroaryl groups. Examples of reduced heteroaryl
25 groups thus include pyrrolidinyl, tetrahydrofuryl, tetrahydrothienyl and piperidinyl.

The compounds of the first aspect of the invention are useful inhibitors of enzymes involved in the degradation of fats. Preferably therefore the first aspect of the invention provides the use of a compound of formula (I) as defined hereinabove, or a
30 pharmaceutically acceptable salt, ester, amide or prodrug thereof, in the manufacture

of a medicament for the control or treatment of obesity, or obesity-related disorders or for promoting non-medical weight loss.

5 Preferably, a compound for use according to the first aspect of the invention is a compound of formula (II).



(II)

10 or a pharmaceutically acceptable salt, ester, amide or prodrug thereof;

wherein:-

R¹, R⁴, R⁵, R⁶, R⁷, X¹ and X² are as defined above for formula I;

15

R² is hydrogen or is a group as defined above for R¹; and

R⁸, R⁹, R¹⁰, R¹¹ are each independently hydrogen, halo, hydroxy, amino, nitro, cyano,

20 or a group R¹, as defined above;

or a group R¹²Q where Q is O, CO, CONH, NHCO, S, SO, SO₂, or SO₂NH₂ and R¹² is hydrogen or a group R¹ as defined above;

or a group R^1R^2N where R^1 and R^2 are as defined above, with the proviso that any hetero atom substituent in R^1 and/or R^2 must be separated from the aromatic hetero atom substituent by at least two carbon atoms (preferably saturated).

5 In the compounds of formula (II):

R^1 preferably represents phenyl substituted by a group selected from OR^{13} , $-COR^{13}$, CO_2R^{13} , SOR^{13} , SO_2R^{13} , $CONR^{13}R^{14}$, $NR^{14}C(O)NR^{13}$, $C_{1-10}alkyl$, $C_{1-10}alkoxy$, halo $C_{1-10}alkyl$, aryl, aryl $C_{1-10}alkyl$, heteroaryl or heteroaryl $C_{1-10}alkyl$; wherein R^{13} and R^{14} each independently represents hydrogen, $C_{1-10}alkyl$, $C_{2-10}alkenyl$, $C_{2-10}alkynyl$, $C_{3-6}cycloalkyl$, $C_{3-6}cycloalkenyl$, aryl, aryl $C_{1-10}alkyl$, heteroaryl, heteroaryl $C_{1-10}alkyl$, reduced heteroaryl or reduced heteroaryl $C_{1-10}alkyl$.

15 More preferably R^1 represents phenyl substituted by OR^{13} or COR^{13} wherein R^{13} is preferably aryl, most preferably phenyl; phenyl substituted by $-CO_2R^{13}$ wherein R^{13} represents $C_{1-10}alkyl$, preferably $C_{1-6}alkyl$; or phenyl substituted by $C_{6-10}alkyl$.

R^2 preferably represents hydrogen or $C_{1-10}alkyl$;

20 R^8 , R^9 , R^{10} and R^{11} preferably each independently represents hydrogen, halo, hydroxy, amino, nitro, cyano, thiol, $C_{1-10}alkyl$, $C_{1-10}alkoxy$, $C_{1-10}cycloalkyl$, $C_{1-10}cycloalkoxy$, $C(O)R^{15}$, $C(O)NR^{15}R^{16}$, $S(O)R^{15}$ or halo $C_{1-10}alkyl$;

25 where R^{15} and R^{16} each independently represent hydrogen or $C_{1-10}alkyl$.

R^8 is hydrogen or halogen e.g. fluorine; most preferably hydrogen;

R^9 is preferably hydrogen or lower branched or unbranched alkyl having 1 to 10 carbon atoms: cyclic alkyl having 3 to 6 carbon atoms, e.g. cyclopropyl; haloC₁₋₆alkyl, e.g. trifluoromethyl; or a halogen, e.g. chlorine or fluorine;

5 R^{10} is preferably hydrogen, lower branched or unbranched alkyl having 1 to 10 carbon atoms e.g. ethyl, butyl or octyl; cyclic alkyl having 3 to 6 carbon atoms, e.g. cyclopropyl; haloC₁₋₆alkyl, e.g. trifluoromethyl or a halogen, e.g. chlorine or fluorine;

10 R^{11} is preferably hydrogen, halogen, eg. fluorine; or branched or unbranched alkyl having 1 to 10 carbon atoms.

15 Preferably, in compounds of formula (II) at least one of R^8 , R^9 , R^{10} and R^{11} represents a substituent other than hydrogen. Thus, for example, R^8 may represent a hydrogen atom and R^9 , R^{10} and R^{11} are as defined above. In a preferred embodiment each of R^8 and R^{11} represents a hydrogen atom, and one or both of R^9 and R^{10} represents a substituent as defined above.

20 Preferably, a compound for use according to the first aspect of the invention comprises a compound of formula (II) or a pharmaceutically acceptable salt, ester, amide or prodrug thereof; wherein:

25 R^1 is aryl e.g. optionally substituted phenyl or 2-naphthyl, or an aryl alkyl group wherein the alkyl moiety has up to 25 e.g. up to 20 carbon atoms, or an aryl aryl group; wherein the aryl alkyl group or the aryl aryl group may be separated by a spacer where the spacer can be an ester, amide, O, CH₂, or a ketone and wherein any aryl group is preferably a phenyl, optionally substituted with alkyl, haloalkyl or halogen;

30 R^2 is hydrogen or is a group as defined above for R^1 ;

R⁸ is hydrogen or fluorine;


R⁹ is lower branched or unbranched alkyl having 1 to 10 carbon atoms; cyclic alkyl having 3 to 10 carbon atoms, e.g. cyclopropyl; haloalkyl, e.g. trifluoromethyl; or a halogen, e.g. chlorine or fluorine;

R¹⁰ is lower branched or unbranched alkyl having 1 to 10 carbon atoms; e.g. ethyl, butyl or octyl, cyclic alkyl having 3 to 10 carbon atoms, e.g. cyclopropyl; haloalkyl, e.g. trifluoromethyl; or a halogen, e.g. chlorine or fluorine;

R¹¹ is hydrogen, lower branched or unbranched alkyl having 1 to 10 carbon atoms, or halogen, e.g. fluorine.

Most preferably, R¹ is unsubstituted phenyl or phenyl substituted by a group selected from C₁₋₈ alkyl, eg butyl, pentyl, hexyl or heptyl; halo-C₁₋₈ alkyl, eg CF₃; OR⁶ where R⁶ is phenyl or COR⁴ where R⁴ is phenyl or C₁₋₈ alkyl.

In a second aspect the present invention provides novel compounds of formula (IIa):



(IIa)

or a pharmaceutically acceptable salt, ester, amide or prodrug thereof;

wherein:-

R^{1a} represents

(i) a C_{10-30} branched or unbranched alkyl, C_{2-30} alkenyl, C_{2-30} alkynyl, cycloalkenyl, ,
 aryl- C_{10-30} alkyl, aryl- C_{10-30} alkenyl, heteroaryl, heteroaryl- C_{1-30} alkyl, heteroaryl- C_{2-30}
 alkenyl, reduced aryl, reduced heteroaryl, reduced heteroaryl- C_{1-30} alkyl or a
 5 substituted derivative thereof wherein the substituents are one or more independently
 chosen from the group consisting of halogen, C_{1-10} alkyl, halosubstituted C_{1-10} alkyl,
 aryl, aryl- C_{1-10} alkyl, heteroaryl, reduced heteroaryl, reduced heteroaryl- C_{1-10} alkyl,
 aryl- C_{1-10} alkoxy, cyano, nitro, $-C(O)R^{13}$, $-CO_2R^{13}$, $-SOR^{13}$, $-SO_2R^{13}$, $-NR^{13}R^{14}$, -
 OR^{13} , $-SR^{13}$, $-C(O)NR^{13}R^{14}$, and $-NR^{14}C(O)R^{13}$, with the proviso that any hetero atom
 10 substituent in R^1 must be separated from the exocyclic nitrogen atom by at least two
 carbon atoms (preferably saturated); or

(ii) aryl substituted by one or more independently chosen from the group consisting of
 halosubstituted C_{1-10} alkyl, aryl, aryl- C_{1-10} alkyl, heteroaryl, reduced heteroaryl,
 15 reduced heteroaryl- C_{1-10} alkyl, aryl- C_{1-10} alkoxy, cyano, $-C(O)R^{13}$, $-CO_2R^{13}$, $-SOR^{13}$, -
 SO_2R^{13} , $-NR^{13}R^{14}$, $-OR^{13}$ (providing that in this instance R^{13} does not represent aryl
 or alkyl), $-SR^{13}$, $-C(O)NR^{13}R^{14}$, and $-NR^{14}C(O)R^{13}$

wherein:-

20 R^{13} and R^{14} each independently represents hydrogen, C_{1-10} alkyl, C_{2-10} alkenyl,
 C_{2-10} alkynyl, C_{3-6} cycloalkyl,
 C_{3-6} cycloalkenyl, aryl, aryl- C_{1-10} alkyl, heteroaryl, heteroaryl- C_{1-10} alkyl, reduced
 heteroaryl or reduced heteroaryl- C_{1-10} alkyl;

25 R^{2a} is hydrogen or is a group as defined above for R^1 ; and

R^{8a} , R^{9a} , R^{10a} , R^{11a} are as defined above for formula (II).

30 provided that:

when R^1 represents a heteroaryl group it is not thiadiazolyl, triazolyl or thiazolyl and when R^1 represents a reduced heteroaryl group it is not thiazolidinyl.

In the compounds of formula (IIa):

5

R^{1a} preferably represents phenyl substituted by a group selected from OR^{13} (providing in this instance R^{13} does not represent alkyl or aryl), $-COR^{13}$, CO_2R^{13} , SOR^{13} , SO_2R^{13} , $CONR^{13}R^{14}$, $NR^{14}C(O)NR^{13}$, $haloC_{1-10}alkyl$, aryl, aryl $C_{1-10}alkyl$, heteroaryl or heteroaryl $C_{1-10}alkyl$.

10

More preferably R^{1a} represents phenyl substituted by COR^{13} wherein R^{13} is preferably aryl, most preferably phenyl; or phenyl substituted by $-CO_2R^{13}$ wherein R^{13} represents $C_{1-10}alkyl$, preferably $C_{1-6}alkyl$.

15

R^{2a} preferably represents hydrogen or $C_{1-10}alkyl$;

R^{8a} , R^{9a} , R^{10a} and R^{11a} preferably each independently represents hydrogen, halo, hydroxy, amino, nitro, cyano, thiol, $C_{1-10}alkyl$, $C_{1-10}alkoxy$, $C_{1-10}cycloalkyl$, $C_{1-10}cycloalkoxy$, $C(O)R^{15}$, $C(O)NR^{15}R^{16}$, $S(O)R^{4a}$ or $haloC_{1-10}alkyl$;

20

where R^{15} and R^{16} each independently represent hydrogen or $C_{1-10}alkyl$.

R^{8a} is hydrogen or halogen e.g. fluorine; most preferably hydrogen;

25

R^{9a} is preferably hydrogen or lower branched or unbranched alkyl having 1 to 10 carbon atoms; cyclic alkyl having 3 to 6 carbon atoms; e.g. cyclopropyl, $haloC_{1-6}alkyl$, e.g. trifluoromethyl or a halogen, e.g. chlorine or fluorine;


R^{10a} is more preferably hydrogen, lower branched or unbranched alkyl having 1 to 10 carbon atoms e.g. ethyl, butyl or octyl; cyclic alkyl having 3 to 6 carbon atoms e.g. cyclopropyl, haloC₁₋₆alkyl e.g. trifluoromethyl or a halogen e.g. chlorine or fluorine;

- 5 R^{11a} is preferably hydrogen, halogen, eg. fluorine; or branched or unbranched alkyl having 1 to 10 carbon atoms.

Preferably, in compounds of formula (IIa) at least one of R^{8a} , R^{9a} , R^{10a} and R^{11a} represents a substituent other than hydrogen. Thus, for example, R^{8a} may represent a hydrogen atom and R^{9a} , R^{10a} and R^{11a} are as defined above. In a preferred
10 embodiment each of R^{8a} and R^{11a} represents a hydrogen atom, and one or both of R^{9a} and R^{10a} represents a substituent as defined above.

In a yet further embodiment the present invention provides compounds of formula (IIb)

15



20

wherein

R^8 - R^{11} are as defined hereinbefore and

- 25 R^{20} represents C₁₋₂₀alkyl, C₁₋₂₀alkoxy, or optionally substituted phenoxy.

Preferred substituents for phenoxy include one or more of halo, CF₃, lower alkyl and lower alkoxy groups.

When R²⁰ represents an alkyl or alkoxy group this preferably contains from 6-12 carbon atoms.

In this embodiment R²⁰ is most preferably phenoxy.

5

Preferred values of R⁸ – R¹¹ are as defined above.

Compounds of formula (IIb) represent a novel selection on the basis of their advantageous activity as lipase inhibitors.

10

Examples of pharmaceutically acceptable salts of the formula include those derived from organic acids such as methanesulphonic acid, benzenesulphonic acid and p-toluenesulphonic acid, mineral acids such as hydrochloric and sulphuric acid and the like, giving methanesulphonate, benzenesulphonate, p-toluenesulphonate,

15 hydrochloride and sulphate, and the like, respectively or those derived from bases such as organic and inorganic bases. Examples of suitable inorganic bases for the formation of salts of compounds for this invention include the hydroxides, carbonates, and bicarbonates of ammonia, lithium, sodium, calcium, potassium, aluminium, magnesium, zinc and the like. Salts can also be formed with suitable organic bases.

20 Such bases suitable for the formation of pharmaceutically acceptable base addition salts with compounds of the present invention include organic bases which are nontoxic and strong enough to form salts. Such organic bases are already well known in the art and may include amino acids such as arginine and lysine, mono-, di-, or trihydroxyalkylamines such as mono-, di-, and triethanolamine, choline, mono-, di-,
25 and trialkylamines, such as methylamine, dimethylamine, and trimethylamine, guanidine; N-methylglucosamine; N-methylpiperazine; morpholine; ethylenediamine; N-benzylphenethylamine; tris(hydroxymethyl) aminomethane; and the like.



Salts may be prepared in a conventional manner using methods well known in the art.
30 Acid addition salts of said basic compounds may be prepared by dissolving the free








base compounds according to the first or second aspects of the invention in aqueous or aqueous alcohol solution or other suitable solvents containing the required acid. Where a compound of formula (I) contains an acidic function a base salt of said compound may be prepared by reacting said compound with a suitable base. The acid or base salt may separate directly or can be obtained by concentrating the solution eg. by evaporation. The compounds of this invention may also exist in solvated or hydrated forms.








The invention also extends to prodrugs of the aforementioned compounds. A prodrug is commonly described as an inactive or protected derivative of an active ingredient or a drug which is converted to the active ingredient or drug in the body.









Representative compounds according to the first and /or second aspects of the invention are those which include;






Table 1







Reference Number	Structure	Compound Name
1		2-Phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
2		2-(4-Butylphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one







3		6-Chloro-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
4		2-Butylamino-4 <i>H</i> -3,1-benzoxazin-4-one
5		6-Methyl-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
6		2-(4-Methoxyphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
7		2-(4-Methylphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
8		2-(4-Phenoxyphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
9		2-(4-Chlorophenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one








10		2-[4(1-Methylethyl)phenylamino]-4 <i>H</i> -3,1-benzoxazin-4-one
11		2-(4-Trifluoromethylphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
12		2-(3-Trifluoromethylphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
13		6-Methyl-2-(naphth-2-ylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
14		2-(4-Butoxycarbonylphenylamino)-6-methyl-4 <i>H</i> -3,1-benzoxazin-4-one
15		6-Methyl-2-(4-phenoxyphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
16		2-Ethylamino-4 <i>H</i> -3,1-benzoxazin-4-one







17		7-Fluoro-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
18		5-Fluoro-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
19		7-Methyl-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
20		7-Ethyl-2-Phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
21		2-(4-Hexylphenylamino)-6-methyl-4 <i>H</i> -3,1-benzoxazin-4-one
22		2-(4-Heptyloxyphenylamino)-6-methyl-4 <i>H</i> -3,1-benzoxazin-4-one
23		7-Octyl-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
24		7-Methyl-2-(4-phenoxyphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one








25		2-Hexadecylamino-6-methyl-4 <i>H</i> -3,1-benzoxazin-4-one
26		7-Butyl-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
27		7-Methyl-2-(2-phenoxyphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
28		7-Methyl-2-(3-phenoxyphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
29		2-(4-Benzoylphenylamino)-7-methyl-4 <i>H</i> -3,1-benzoxazin-4-one








30		2-(4-Phenoxyphenylamino)-7-trifluoromethyl-4H-3,1-benzoxazin-4-one
31		7-Methyl-2-(4-octylphenylamino)-4H-3,1-benzoxazin-4-one
32		2-Phenylamino-4H-pyrido[3,4-d][1,3]oxazin-4-one
33		2-(2-Cyanophenylamino)-7-methyl-4H-3,1-benzoxazin-4-one
34		6-Nitro-2-phenylamino-4H-3,1-benzoxazin-4-one
35		6-Acetamido-2-phenylamino-4H-3,1-benzoxazin-4-one



36		2-Phenylamino-7-trifluoromethyl-4H-3,1-benzoxazin-4-one
37		7-Amino-2-phenylamino-4H-3,1-benzoxazin-4-one
38		2-Phenylamino-4H-pyrido[2,3-d][1,3]oxazin-4-one
39		2-Cyclopropylamino-4H-3,1-benzoxazin-4-one
40		2-(3-Cyanophenylamino)-7-methyl-4H-3,1-benzoxazin-4-one
41		2-(4-Cyanophenylamino)-4H-3,1-benzoxazin-4-one

42		2-(4-Cyanophenylamino)-7-methyl-4 <i>H</i> -3,1-benzoxazin-4-one
43		2-(4-Carboxyphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
44		2-(4-Aminophenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
45		2-(4-Hydroxyphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
46		2-(4- <i>N</i> -Methylcarbamoylphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
47		2,2'-(1,8-Octylenediamino)-bis-4 <i>H</i> -3,1-benzoxazin-4-one
48		2-(2-Phenoxyphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one

49		2-(3-phenoxyphenylamino)- 4 <i>H</i> -3,1-benzoxazin-4-one
50		2-(Naphth-2-ylamino)-4 <i>H</i> - 3,1-benzoxazin-4-one
51		2-(6-Phenylhexylamino)-4 <i>H</i> - 3,1-benzoxazin-4-one
52		2-(Pyrrol-3-ylamino)-4 <i>H</i> -3,1- benzoxazin-4-one
53		2-(Piperidin-4-ylamino)-4 <i>H</i> - 3,1-benzoxazin-4-one
54		2-[6-(Pyrrol-2-yl)- hexylamino]-4 <i>H</i> -3,1- benzoxazin-4-one

55		2-(4-Ethoxycarbonylphenylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
56		6-Cyano-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
57		2-Phenyl-6-Trifluoromethyl-4 <i>H</i> -3,1-benzoxazin-4-one
58		6-Formyl-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
59		2-Phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one-6-sulfinic acid
60		7-Hydroxy-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
61		7-Cyclopropyl-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one

62		6,7-Dimethyl-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
63		6-Iodo-2-octylamino-4 <i>H</i> -3,1-benzoxazin-4-one
64		7-Butyl-2-octylamino-4 <i>H</i> -3,1-benzoxazin-4-one
65		6-Methyl-2-(dodeca-7-ynylamino)-4 <i>H</i> -3,1-benzoxazin-4-one
66		6-Methyl-2-[6-(thien-2-yl)hexylamino]-4 <i>H</i> -3,1-benzoxazin-4-one
67		8-Fluoro-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one
68		6-Cyclopropyl-2-phenylamino-4 <i>H</i> -3,1-benzoxazin-4-one

69		6-Mercapto-2-phenylamino- <i>4H</i> -3,1-benzoxazin-4-one
70		6-Cyano-2-phenylamino- <i>4H</i> -3,1-benzoxazin-4-one

Compounds 2, 3, 5, 6, 8, 11-15 and 17-70 in Table 1 above are believed to be novel and as such represent preferred embodiments of the present invention.

- 5 Preferred compounds of formula (II) listed in Table 1 include compounds 1, 3, 5, 9, 17, 19, 20, 23 and 26.

Preferred compounds of formula (IIa) listed in Table 1 include compounds 11, 12, 14, 25, 29 and 30.

10

Preferred compounds of formula (IIb) listed in Table 1 include compounds 2, 6, 7, 8, 10, 15, 21, 24.

Particularly preferred compounds of formula (IIa) and (IIb) are:

- 15 2-(4-Phenoxyphenylamino)-*4H*-3,1-benzoxazin-4-one
 2-(4-Butoxycarbonylphenylamino)-6-methyl-*4H*-3,1-benzoxazin-4-one
 6-Methyl-2-(4-phenoxyphenylamino)-*4H*-3,1-benzoxazin-4-one
 2-(4-Hexylphenylamino)-6-methyl-*4H*-3,1-benzoxazin-4-one
 7-Methyl-2-(4-phenoxyphenylamino)-*4H*-3,1-benzoxazin-4-one
 20 2-(4-Benzoylphenylamino)-7-methyl-*4H*-3,1-benzoxazin-4-one
 2-(4-Phenoxyphenylamino)-7-trifluoromethyl-*4H*-3,1-benzoxazin-4-one


Preferred compounds of the invention listed above extend to the tautomers thereof, as well as (but not limited to) pharmaceutically acceptable salts, esters, amides or prodrugs thereof or a derivative with one or more lipid groups (natural or synthetic) attached.

5

A third aspect of the invention provides a process for the manufacture of any one or more of the novel compounds or derivatives according to the first and/or second aspects of the invention. Thus, the present invention provides a process for the preparation of a novel compound of formula (II) in particular a compound of formula (IIa) which process comprises:

10

Process (A) cyclising a compound of formula (III)




15

wherein R¹ and R⁸-R¹¹ are as hereinbefore defined and R¹⁸ is hydrogen or C₁₋₆alkyl.

20

or:

Process (B) reacting a compound of formula (IV)



25

30

with an amine of formula (V)



or:

Process (C) converting a compound of formula (I), (II), (IIa) or (IIb) into a different compound of formula (IIa) or (IIb), by, for example,

(i) reduction of a compound of formula (I), (II), (IIa) or (IIb) wherein any of R^1 , R^8 , R^9 , R^{10} and R^{11} contains an alkenyl or alkynyl group or moiety, to the corresponding alkyl or alkenyl group or moiety; or

(ii) alkylation of a compound of formula (I), (II), (IIa) or (IIb) where one or more of R^8 , R^9 , R^{10} and R^{11} represents a halogen atom.

Process (A) may be effected by reacting a compound (III) with a dehydrating agent in an organic solvent. Suitable dehydrating agents include sulphuric acid, and when R^{18} is hydrogen, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) or polymer supported EDC. The reaction may be effected at a temperature in the range 10 to 50°C, preferably ambient temperature e.g. 20-30°C. When polymer supported EDC is employed it may be removed by filtration at the end of the reaction, and the product isolated from solution by standard procedures, such as removal of the solvent and purification by flash column chromatography. Alternatively the cyclisation may be effected using concentrated sulphuric acid.


Alternatively, cyclisation according to process (A) may be effected by reaction with excess chloroformate or by addition of another cyclisation reagent, which promotes ring closure. Suitable cyclisation reagents include for example, methyl chloroformate, carbonyl diimidazole, acetic anhydride, phosgene, oxalyl chloride, thionyl chloride or

a peptide coupling agent such as dicyclohexyl carbodiimide (DCC). The cyclisation reagent is preferably phosgene, triphosgene or thionyl chloride. When a chloroformate is employed this is preferably a low molecular weight chloroformate, on grounds of cost and ease of removing the resulting alcohol.

5

Compounds of the formula (III) may themselves be prepared according to a variety of methods. Thus for example a compound of formula:

10



(VI)

15

can be reacted with an isocyanate of formula (VII):




20

The reaction is preferably carried out in an inert organic solvent, such as an ether e.g. tetrahydrofuran, an aliphatic hydrocarbon such as pentane or hexane; a halogenated hydrocarbon such as dichloromethane; or an aromatic hydrocarbon such as benzene or toluene, and usually at ambient temperature. The intermediate urea may cyclise directly in a 'one pot' reaction, without isolation. Alternatively, if desired the urea may be isolated prior to cyclisation. Similarly any unreacted urea intermediate may be cyclised in a subsequent reaction step. It will be appreciated that the above reaction results in a compound (III) where R² is hydrogen.

25


Alternatively a compound of formula (III) may be prepared by reacting an isocyanate of formula (VIII):



(wherein R⁸, R⁹, R¹⁰, R¹¹ and R¹⁸ are as hereinbefore defined)

10 with an amine of formula (V) R¹R²NH.

Compounds of formula (III) may also be prepared from compounds of formula (IX)



20 by reaction with an amine R¹R²NH.

Compounds (IX) may themselves be prepared by reacting a compound (VI) with an amine (V) in the presence of trichloromethyl chloroformate and in a solvent such as tetrahydrofuran or dimethyl formamide.

25

Process (B) may be effected by reacting a compound of formula (IV) with an amine R¹R²NH in the presence of a base e.g. sodium hydroxide, followed by cyclisation, for example as described for process (A).

30 Compounds of formula (IV) may be obtained by cyclisation of a compound of formula (VI) wherein R¹⁸ is hydrogen using for example phosgene or a synthetic equivalent.

In process (C), reduction of an alkenyl or alkynyl group may be effected for example by catalytic hydrogenation using e.g. 10% palladium on charcoal in an alcoholic solvent, such as ethanol, under 1 atmosphere of hydrogen gas.

5 Alkylation according to process (C)(ii) may be effected using a Stille or other palladium catalysed cross-coupling process, using e.g. tetra-alkyl tin such as tetramethyl tin and $\text{PhCH}_2\text{Pd}(\text{PPh}_3)_2\text{Cl}$ in HMPA at elevated temperature e.g. 50-100°C. Other halides or pseudohalides e.g. triflates may be employed as starting materials.

10

Further methodology for preparing 2-amino-1,3-benzoxazin-4-one derivatives is described in *J. Med. Chem.* 1990, **33**(2):464-479 and *J. Med. Chem.* 1998 **41**:1060-1067, as well as US Patent No. 4,657,893.

15 A fourth aspect of the invention is a compound according to the first and/or second aspects of the invention (i.e. compounds of formulae (I), (II) and (IIa)), for use in medicine. Preferred features of the first and second aspects of the invention also apply to the fourth aspect. Further details of the fourth aspect of the invention are set out in the text which follows.

20

A fifth aspect of the invention relates to a compound according to the first and/or second aspects of the invention for use in the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality. This includes both *in vivo* and *in vitro* uses and other uses such as industrial uses. Such an enzyme is one which catalyses the breakdown of a substrate containing an ester functionality by the addition of water, resulting in the cleavage of a chemical bond. Such enzymes are involved in key processes in the body. Enzymes according to this invention include lipases (hydrolyse fatty acid esters), esterases (hydrolyse esters) and phosphatases (hydrolyse phosphate esters).

25
30

The enzyme is preferably a lipase. Lipases include pancreatic lipase, gastric lipase, lipoprotein lipase, lingual lipase, adipose tissue lipase, hormone sensitive lipase, phospholipase A1, A2, B, C, D etc., hepatic lipase, and other triacyl, diacyl and monoacylglycerol lipases in the mammalian body. Many similar such lipases are also
5 known in plants, fungi and microorganisms.

Also covered are esterase enzymes and phosphatase enzymes. Esterase enzymes include pig liver esterase, cholesteryl esterase, retinyl esterase, 1-alkyl-2-glycerophosphocholine esterase, carboxylic ester hydrolases and cholesterol esterase.
10 Phosphatase enzymes include serine/threonine phosphatases PP1, PP2 and PP3, phosphoprotein phosphatase, myosin-light-chain phosphatase, protein phosphatase 2C and protein tyrosine phosphatase.

Compounds according to the invention, for use in medicine, are primarily for use in
15 relation to the prevention and/or treatment of a medical condition such as obesity, hyperlipaemia, hyperlipidaemia and related diseases such as hyperglycaemia (type II diabetes), hypertension, cardiovascular disease, stroke, gastrointestinal disease and gastrointestinal conditions. Compounds according to the first and/or second aspects of the invention are useful in these and other conditions due to their ability to inhibit an
20 enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality. The invention also relates to non-medical weight loss, such as cosmetic weight loss and includes improving bodily appearance in general. Throughout this text, the prevention and/or treatment of any disorder means any effect which mitigates any damage or any medical disorder, to any extent, and includes prevention and
25 treatment themselves. The term "treatment" means any amelioration of disorder, disease, syndrome, condition, pain or a combination of two or more thereof.

Clearly, an important application of the invention is in relation to weight loss (of all kinds as described above) in humans. However, the invention applies to medical and
30 non-medical weight loss in any animal whose metabolism of fat and fat derivatives

involves an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality. Thus, the invention has veterinary application and is particularly useful in relation to medical and non-medical weight loss in companion animals such as pet cats and dogs as well as in animals which provide meat for human consumption.

5 In the case of the latter, the application of the present invention is to reduce fat content in order to provide a leaner meat product.

It is also believed that the compounds may be useful in reducing levels of toxins (e.g. dioxins and PCBs) stored in body fat. Without wishing to be bound by theory, it is
10 believed that increasing the amount of undigested fat passing through the body enhances diffusion of toxins from fat stored in the body into fats in the blood, and thence into the intestine.

The fifth aspect of the invention has important applications. It includes test and
15 diagnostic methods and the control and inhibition of unwanted enzymes, preferably lipases, in any process or in any product. The processes or products which preferably involve a lipase include processing of agricultural commodities (e.g. oilseeds), recovery and isolation of enzymes from biotechnological processes (e.g. involving lysis of microorganisms), the manufacture and extraction of crude oil (especially oil
20 and plastics), the industrial manufacture of triglycerides or other fats, manufacture of healthcare goods which comprise surfactants, soap or detergent (e.g. bath oils, creams), the manufacturing and processing of liposomes (e.g. healthcare products, diagnostics, gene therapy), the treatment of industrial waste (e.g. paper effluent treatment) and preventing the degradation of foodstuff which comprises a fat (e.g.
25 chocolate processing). Thus, the invention also relates to these products and processes, e.g. a foodstuff which comprises a compound according to the first aspect of the invention, in particular foodstuffs which have a high fat content such as cakes, biscuits, pastry-products and the like and chocolate products. The preferred features of the fifth aspect of the invention, including the preferred enzymes are as discussed
30 for the previous aspects of the invention.

A sixth aspect of the invention provides a composition comprising a novel compound according to the first and second aspects of the invention, in combination with a pharmaceutically acceptable carrier or diluent. Suitable carriers and/or diluents are well known in the art and include pharmaceutical grade starch, mannitol, lactose, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose (or other sugar), magnesium carbonate, gelatin, oil, alcohol, detergents, emulsifiers or water (preferably sterile). The composition may be a mixed preparation of a composition or may be a combined preparation for simultaneous, separate or sequential use (including administration).

The compounds according to the invention for use in the aforementioned indications may be administered by any convenient method, for example by oral (including by inhalation), parenteral, mucosal (e.g. buccal, sublingual, nasal), rectal or transdermal administration and the compositions adapted accordingly.

For oral administration, the compounds can be formulated as liquids or solids, for example solutions, syrups, suspensions or emulsions, tablets, capsules and lozenges.

A liquid formulation will generally consist of a suspension or solution of the compound or physiologically acceptable salt in a suitable aqueous or non-aqueous liquid carrier(s) for example water, ethanol, glycerine, polyethylene glycol or an oil. The formulation may also contain a suspending agent, preservative, flavouring or colouring agent.

A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and microcrystalline cellulose.

A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, powders, granules or pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable
5 pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule.

Compositions for oral administration may be designed to protect the active ingredient against degradation as it passes through the alimentary tract, for example by an outer
10 coating of the formulation on a tablet or capsule.

Typical parenteral compositions consist of a solution or suspension of the compound or physiologically acceptable salt in a sterile aqueous carrier or non-aqueous or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and
15 then reconstituted with a suitable solvent just prior to administration.

Compositions for nasal or oral administration may conveniently be formulated as aerosols, drops, gels and powders. Aerosol formulations typically comprise a solution
20 or fine suspension of the active substance in a physiologically acceptable aqueous or non-aqueous solvent and are usually presented in single or multidose quantities in sterile form in a sealed container, which can take the form of a cartridge or refill for use with an atomising device. Alternatively the sealed container may be a unitary dispensing device such as a single dose nasal inhaler or an aerosol dispenser fitted
25 with a metering valve which is intended for disposal once the contents of the container have been exhausted. Where the dosage form comprises an aerosol dispenser, it will contain a pharmaceutically acceptable propellant. . The aerosol dosage forms can also take the form of a pump-atomiser.

Compositions suitable for buccal or sublingual administration include tablets, lozenges and pastilles, wherein the active ingredient is formulated with a carrier such as sugar and acacia, tragacanth, or gelatin and glycerin.

- 5 Compositions for rectal or vaginal administration are conveniently in the form of suppositories (containing a conventional suppository base such as cocoa butter), pessaries, vaginal tabs, foams or enemas.

- 10 Compositions suitable for transdermal administration include ointments, gels and patches, and injections, including powder injections.

Conveniently the composition is in unit dose form such as a tablet, capsule or ampoule.

- 15 The compositions of the sixth aspect of the invention are useful in the prevention and/or treatment of obesity, obesity-related disorder, other medical weight loss and non-medical related weight loss. Preferred features of this aspect of the invention are as described above for the first to fifth aspects of the invention.

- 20 A seventh aspect of the invention provides a process for the manufacture of a composition according to the sixth aspect of the invention. The manufacture can be carried out by standard techniques well known in the art and involves combining a compound according to the first or second aspect of the invention and the pharmaceutically acceptable carrier or diluent. The composition may be in any form
25 including a tablet, a liquid, a capsule, and a powder or in the form of a food product, e.g. a functional food. In the latter case the food product itself may act as the pharmaceutically acceptable carrier.

- 30 An eighth aspect of the invention provides a method for the prevention and/or treatment of obesity or an obesity-related disorder, the method comprising the

administration of a compound according to the first or second aspect of the invention, preferably in combination with a pharmaceutically acceptable carrier or diluent (as per the sixth aspect of the invention). Obesity-related disorders include hyperlipemia, hyperlipideamia, hyperglycaemia, hypertension, cardiovascular disease, stroke, gastrointestinal disease and gastrointestinal conditions. The compound or composition is preferably administered to a patient in need thereof and in a quantity sufficient to prevent and/or treat the symptoms of the condition, disorder or disease. For all aspects of the invention, particularly medical ones, the administration of a compound or composition has a dosage regime which will ultimately be determined by the attending physician and will take into consideration such factors such as the compound being used, animal type, age, weight, severity of symptoms, method of administration, adverse reactions and/or other contraindications. Specific defined dosage ranges can be determined by standard design clinical trials with patient progress and recovery being fully monitored. Such trials may use an escalating dose design using a low percentage of the maximum tolerated dose in animals as the starting dose in man.

The physiologically acceptable compounds of the invention will normally be administered in a daily dosage regimen (for an adult patient) of, for example, an oral dose of between 1 mg and 2000 mg, preferably between 30 mg and 1000 mg, e.g. between 10 and 250 mg or an intravenous, subcutaneous, or intramuscular dose of between 0.1 mg and 100 mg, preferably between 0.1 mg and 50 mg, e.g. between 1 and 25 mg of the compound of the formula (I) or a physiologically acceptable salt thereof calculated as the free base, the compound being administered 1 to 4 times per day. Suitably the compounds will be administered for a period of continuous therapy, for example for a week or more.

A ninth aspect of the invention provides a cosmetic method for maintaining a given weight, or for cosmetic weight loss, the method comprising the administration of a compound according to the first aspect of the invention, preferably in combination with a pharmaceutically acceptable carrier or diluent (as per the fifth aspect of the

invention). The "medicament" is preferably administered to a patient in need thereof and in a quantity sufficient to maintain a given weight or for cosmetic weight loss.

The eighth and ninth aspects of the invention relate to methods of treatment of humans and other animals, in particular companion animals and other animals which provide meat for human consumption, such as cattle, pigs and sheep (all of any age).

The invention will now be described with reference to the following non-limiting examples.

Biological Test Methods and Results

Test Compounds

The benzoxazinone compounds used in the following tests are identified by the reference number assigned in Table 1 hereinbefore.

Measurement of lipase activity using a quinine diimine dye colorimetric assay

The inhibitory activity of the selected compounds to pancreatic lipase was measured in the following assay available from Sigma Ltd (Lipase PSTM, catalog number 805-A):


Pancreatic lipase

1,2-dibutylin ----- > 2-monoglyceride + fatty acid


Monoglyceride lipase

2-monoglyceride ----- > glycerol + fatty acid


Glycerol kinase



Glycerol phosphate oxidase



Peroxidase



10

The glycerol released from the action of pancreatic and monoglyceride lipase was oxidised to release H_2O_2 . The peroxidase reaction step then produces a quinine dye which is pink in colour and absorbs light at a wavelength of 550 nm.

15 Inhibitor

Individual compounds were dissolved in DMSO (dimethyl sulphoxide) at 10 mM. DMSO was used to avoid any problems with compounds being water-insoluble.

20

For individual compounds, the IC_{50} (concentration at which lipase activity is inhibited to one half of the maximum) was calculated by measuring the inhibitory activity from log-dose response curves using a range of inhibitor concentrations.

Results

25

A range of compounds were assayed in the quinine diimine dye colorimetric assay which provides a rapid method to measure lipase inhibitory activity. None of the compounds tested interfered with the colorimetric reaction, i.e. they did not give false positive results.

A range of inhibitory activities for the tested benzoxazinone compounds were observed, indicating that these compounds are inhibitors of human pancreatic lipase. The following compounds had IC_{50} 's $\leq 1\mu M$: 1-3, 5-12, 14, 15, 17, 19-21, 23-26, 28-30.

5

Measurement of lipase enzyme activity using a NaOH titration method

The inhibitory activity of the selected compounds to pancreatic lipase was measured in the assay described in Pasquier *et al.*, 1986, Vol 7, *Nutritional Biochemistry*, 293-302.

10

Log dose/response curves were constructed using a range of inhibitor concentrations.

Results

15

Selected benzoxazinone compounds were tested in the NaOH titration assay. In this assay, the activity of porcine pancreatic lipase in a system containing lipid micelles is recorded. These conditions are therefore similar to those encountered in the gastrointestinal tract.

20

A range of inhibitory activities were observed for the tested benzoxazinone compounds in this assay indicating that these compounds are inhibitors of porcine pancreatic lipase. The following compounds had an $IC_{50} \leq 2\mu M$: 1-3, 5, 8, 11, 12, 14-20, 24, 26, 28, 29, 30.

25

Thus, the results demonstrate that the tested benzoxazinones are inhibitors of fat digestion and that these compounds may be particularly suitable for the treatment of obesity.

Mouse Model Assay


- 5 Compound 24 was assayed in a mouse model as described by Isler *et al.*, *British Journal of Nutrition*, 1995, 73:851-862 and was found to be a potent lipase inhibitor.

Synthesis of Intermediates

10 Synthesis of 4-substituted anthranilic acids

Example: 4-octyl anthranilic acid (4-octyl-2-aminobenzoic acid)


Method based on that of L.A. Paquette et al. *J.Am.Chem.Soc.* 99, 3734 (1981)




- A solution of 1-bromo-4-octylbenzene (9.9g, 36mmol) in sulfuric acid (20ml) was cooled in an ice bath. To this was added nitric acid (1.44ml, 36mmol). The ice bath was removed and the mixture stirred at room temperature for 20 minutes. A further portion of nitric acid was added (0.07ml, 1.75mmol), stirring being continued for a further 20 min. The mixture was poured into aqueous potassium carbonate, which was extracted with ethyl acetate. The organic extract was washed with saturated aqueous potassium carbonate, water and brine then dried (MgSO₄) and concentrated.
- 20
- Purification of the crude product by flash chromatography (1% EtOAc/hexane)
- 25 removed the unwanted (major) regioisomer and afforded the desired material as a yellow oil (1.7g, 5.4mmol).



5 The substrate (1.7g, 5.4mmol), copper (I) cyanide (0.533g, 5.9mmol) and pyridine (20ml) were refluxed at 150°C for 2 days. Concentration *in vacuo* and purification by flash chromatography (10% to 20% EtOAc/hexane) gave the desired material as a brown oil (739mg, 2.8mmol)



10 The substrate (694mg, 2.7mmol) was heated at 150°C in a mixture of water (2ml), AcOH (1ml) and sulfuric acid (1ml) for 2 days. The mixture was extracted with ethyl acetate, the organic phase being washed with water (x 2), dried (Na₂SO₄) and concentrated to give the desired material (744mg, 2.7mmol).




20 The starting material (744mg, 2.7mmol) was dissolved in ethanol (10ml) and to this was added a slurry of 10% palladium on charcoal (40mg) in ethanol (4ml). The flask was flushed with nitrogen then hydrogen (1 atm) after which stirring was maintained overnight. Further portions of catalyst (5mg and 25mg) were added, the reaction being

complete after a further 24h. The reaction mixture was filtered through celite, thoroughly rinsing with methanol and ethyl acetate. Concentration gave the anthranilic acid (597mg, 2.4mmol) of sufficient purity for use without further purification; δ_{H} (400 MHz, CDCl_3) 0.79-0.81 (3H, m, Me), 1.12-1.36 (10H, m, 5 x CH_2), 1.52 (2H, br.s, ArCH_2CH_2), 2.45 (2H, br.s, ArCH_2), 6.42 (2H, br.s, 2 x ArH), 7.74 (1H, br.s, ArH); m/z (ES^+) 250 (MH^+).

Synthesis of substituted phenyl isocyanates

10 *Example: Preparation of 4-octylphenyl isocyanate*



15 A solution of 4-octylaniline (0.3 ml, 1.3 mmol) and diisopropylamine (0.205 ml, 5.2 mmol) in THF (5 ml) was cooled to -10°C . A 20% solution of phosgene in toluene (1.3 ml, 2.6 mmol) was added, then the mixture was allowed to warm to r.t. and maintained at that temperature for 3h. Excess phosgene was removed under a stream of nitrogen (scrubbed on exit with $\text{NaOH}_{(\text{aq})}$) to give a solution of the crude isocyanate, which was used directly in the next step.

20

4-Benzoylphenylisocyanate was prepared by an analogous procedure from the corresponding aniline.


25 Substituted 4-phenoxyphenyl isocyanates can be prepared from the corresponding amines by known procedures.

Synthesis of compounds according to the invention

Example 1Synthesis of 2-(4-Butoxycarbonylphenylamino)-6-methyl-4H-3,1-benzoxazin-4-one

(Reference number 14)

5



10

A solution of 2-amino-5-methylbenzoic acid (690 mg, 4.57 mmol) in THF (2 ml) was treated with 4-*n*-butoxycarbonylphenyl isocyanate (1.0 g, 4.57 mmol). The mixture was kept at room temperature for 24 h, during which time the solvent was allowed to evaporate to leave a pale brown solid (1.7 g, quant.); δ_H (400 MHz, DMSO- d_6) 0.93 (3H, t, J 7, CH_2CH_3), 1.41 (2H, tq, J, J' 7, CH_2CH_3), 1.67 (2H, tt, J, J' 7, $CH_2CH_2CH_3$), 2.28 (3H, s, CH_3), 4.23 (2H, t, J 7, OCH_2), 7.37 (1H, d, J 8, Ph), 7.77 (1H, s, Ph), 7.85-7.92 (4H, m, Ph), 8.24 (1H, d, J 8, Ph).

15



20


To a solution of the urea (185 mg, 0.5 mmol) in DMF (10 ml) was added polymer-supported EDC (PS-EDC) (0.8 mmol g^{-1} , 1.0 g). The resulting mixture was agitated at room temperature for 18 h, after which the resin was filtered off and washed with DMF (2 x 5 ml). The filtrate and washings were combined and evaporated under reduced pressure to afford the required compound as an off-white solid (150 mg, 85%); δ_H (400 MHz, DMSO- d_6) 0.94 (3H, t, J 7, CH_2CH_3), 1.42 (2H, tq, J, J' 7, CH_2CH_3), 1.69 (2H, tt, J, J' 7, $CH_2CH_2CH_3$), 2.39 (3H, s, CH_3), 4.25 (2H, t, J 7, OCH_2), 7.36 (1H, d, J 8, Ph), 7.63 (1H, d, J 8, Ph), 7.80 (1H, s, Ph), 7.90-7.95 (4H, m, Ph); m/z (ES $^-$) 351 ($M-H^+$).

25

Example 2

Synthesis of 6-Methyl-2-(4-phenoxyphenylamino)-4H-3,1-benzoxazin-4-one


(Reference number 15)



A solution of 2-amino-5-methylbenzoic acid (172 mg, 1.10 mmol) in THF (1 ml) was treated with 4-phenoxyphenyl isocyanate (241 mg, 1.10 mmol). The mixture was kept at room temperature for 24 h, during which time the solvent was allowed to evaporate to leave a pale brown solid. This was dissolved in DMF (5 ml) and added to a suspension of PS-EDC (0.8 mmol g^{-1} , 2.8 g) in DMF (20 ml). The resulting mixture was agitated at room temperature for 18 h, after which the resin was filtered off and washed with DMF (2 x 5 ml). The filtrate and washings were combined and evaporated under reduced pressure. Flash column chromatography over silica (20% ethyl acetate in hexane as eluent) afforded the required compound as an off-white solid (153 mg, 40%); δ_{H} (400 MHz, DMSO- d_6) 2.36 (3H, s, CH_3), 6.96 (2H, d, J 8, Ph), 7.04 (1H, d, J 8, Ph), 7.09 (1H, t, J 8, Ph), 7.26 (1H, d, J 8, Ph), 7.34-7.38 (2H, m, Ph), 7.55-7.56 (1H, m, Ph), 7.76-7.78 (4H, m, Ph); m/z (ES^+) 345 (MH^+).

Example 3

2-(3-chlorophenylamino)-6-octyl-4H-3,1-benzoxazin-4-one




The anthranilic acid (200mg, 0.8 mmol) was dissolved in anhydrous THF (1ml), and to this was added 3-chlorophenylisocyanate (117 μ l, 0.96 mmol). The mixture was stirred for 3h before being concentrated. The residue was partitioned between water and ethyl acetate. The organic layer was washed with 5% aqueous citric acid, saturated aqueous sodium bicarbonate and brine then dried (MgSO₄) and concentrated to give an orange solid. This was purified by flash chromatography on silica (15% EtOAc/hexane to 100% EtOAc, then 10% EtOH/EtOAc) which gave, in order of elution, the benzoxazinone (18mg, 0.05 mmol); δ_H (400 MHz, CDCl₃) 0.77-0.82 (3H, m, Me), 1.19-1.24 (10H, m, 5 x CH₂), 1.55-1.57 (2H, m, ArCH₂CH₂), 2.56-2.62 (2H, m, ArCH₂), 7.00-7.32 (4H, m, 3 x ArH, NH), 7.40-7.49 (2H, m, 2 x ArH), 7.81-7.83 (2H, m, ArH); m/z (ES⁺) 385 (MH⁺) and the urea (160mg, 0.4 mmol); m/z (ES⁻) 401 (M-H).

The residual urea may be cyclised in a separate step (as in the procedure below) to afford more benzoxazinone if required.

Example 4


7-Octyl-2-phenylamino-4H-3,1-benzoxazin-4-one (compound 23)



The urea (126mg, 0.34 mmol) was suspended in dry DCM (4ml). 1-[3-
 5 (Dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (EDC, 131mg, 0.68
 mmol) was added and the mixture stirred for 24h. A further portion of EDC (131mg,
 0.68 mmol) was added and the mixture was stirred for a further 24h. The mixture was
 diluted with ethyl acetate and washed with water, saturated aqueous sodium
 bicarbonate and brine, then dried (Na_2SO_4) and concentrated. This gave the desired
 10 benzoxazinone (89mg, 0.25 mmol), which did not require further purification: δ_{H} (400
 MHz, CDCl_3) 0.92 (3H, t, J 6.7, Me), 1.30-1.36 (10H, m, 5 x CH_2), 1.68-1.70 (2H, m,
 ArCH_2CH_2), 2.71-2.75 (2H, m, ArCH_2), 6.80 (1H, br.s, NH), 7.13-7.21 (2H, m, 2 x
 ArH), 7.28 (1H, s, ArH), 7.42-7.46 (2H, m, ArH), 7.70 (2H, d, J 8.2, ArH), 8.04 (1H,
 d, J 8.1, ArH); m/z (ES^+) 351 (MH^+).


Example 5

7-Methyl-2-(4-phenoxyphenyl)-4H-3,1-benzoxazin-4-one:



2-Amino-4-methylbenzoic acid (207 mg, 1.37 mmol) in THF (3 ml) was treated with
 4-phenoxyphenyl isocyanate (289 mg, 1.37 mmol). The mixture was kept at r.t. for
 48h then diluted with ethyl acetate and washed with 2N HCl, water, saturated aqueous

sodium bicarbonate and brine, dried (Na_2SO_4) and concentrated. The residue was chromatographed on silica (50% to 100% ethyl acetate/hexane gradient, then 1% methanol/ethyl acetate) to give a white solid (217 mg, 0.6 mmol, 44%); δ_{H} (400 MHz, DMSO- d_6) 2.28 (3H, s, Me), 6.72 (1H, d, J 8.0, ArH), 6.95 (4H, m, ArH), 7.08 (1H, t, J 6.9, ArH), 7.36 (2H, t, J 7.9, ArH), 7.56 (2H, d, J 8.7, ArH), 7.87 (1H, d, J 7.6, ArH), 8.11 (1H, s, ArH), 9.49 (1H, s, NH); m/z (ES^-) 362 (M-H^-).



The urea (217 mg, 0.6 mmol) in DCM (4 ml) was treated with EDC (126mg, 0.66 mmol). After 24h and 48h, further portions of EDC were added (115 mg each, 0.6 mmol). The mixture was diluted with ethyl acetate and washed with water, saturated aqueous sodium bicarbonate and brine then dried (Na_2SO_4) and concentrated to afford the benzoxazinone (137 mg, 0.4 mmol, 67%); δ_{H} (400 MHz, CDCl_3) 2.37 (3H, s, Me), 6.7 (1H, br.s, NH), 6.94-7.05 (6H, m, ArH), 7.14 (1H, s, ArH), 7.27 (2H, t, J 7.7, ArH), 7.54 (2H, d, J 8.7, ArH), 7.89 (1H, d, J 8.1, ArH); m/z (ES^+) 345 (MH^+).

The other compounds listed in Table 1 may be prepared in a similar manner to Examples 1 to 5 above. In particular the following compounds were prepared using the starting materials indicated:

Compound number	Starting material 1	Starting material 2
2	2-aminobenzoic acid	4-butylphenyl isocyanate


8	2-aminobenzoic acid	4-phenoxyphenyl isocyanate
10	2-aminobenzoic acid	4(1-methylethyl)phenyl isocyanate
11	2-aminobenzoic acid	4-trifluoromethylphenyl isocyanate
12	2-aminobenzoic acid	3-trifluoromethylphenyl isocyanate
13	2-amino-5-methylbenzoic acid	1-naphthyl isocyanate
14	2-amino-5-methylbenzoic acid	4-butoxycarbonylphenyl isocyanate
15	2-amino-5-methylbenzoic acid	4-phenoxyphenyl isocyanate
17	2-amino-4-fluorobenzoic acid	Phenyl isocyanate
18	2-amino-6-fluorobenzoic acid	Phenyl isocyanate
19	2-amino-4-methylbenzoic acid	Phenyl isocyanate
20	2-amino-4-ethylbenzoic acid	phenyl isocyanate
21	2-amino-5-methylbenzoic acid	4-hexyl phenyl isocyanate
22	2-amino-5-methylbenzoic acid	4-heptyloxyphenyl isocyanate
23	2-amino-4-octylbenzoic acid	Phenyl isocyanate
24	2-amino-4-methylbenzoic acid	4-phenoxyphenyl isocyanate
25	2-amino-5-methylbenzoic acid	Hexadecylisocyanate
26	2-amino-4-butylbenzoic acid	phenyl isocyanate
27	2-amino-4-methylbenzoic acid	2-phenoxyphenylisocyanate
28	2-amino-4-methylbenzoic acid	3-phenoxyphenylisocyanate
29	2-amino-4-methylbenzoic acid	4-benzoylphenyl isocyanate
30	2-amino-4-trifluoromethyl benzoic acid	4-phenoxyphenyl isocyanate
31	2-amino-4-methylbenzoic acid	4-octyl phenyl isocyanate
32	3-aminopyridine-4-carboxylic acid	Phenyl isocyanate
33	2-amino-4-methylbenzoic acid	cyanophenyl isocyanate

The foregoing description details specific compounds, compositions, methods and uses which can be employed to practise the present invention. However, those skilled

in the art will know how to use alternative reliable methods for aiming at alternative embodiments of the invention which are herein encompassed.

CLAIMS

1. The use of a compound comprising formula (I):



(I)

or a pharmaceutically acceptable salt, ester, amide or prodrug thereof; in the
 manufacture of a medicament for the prevention or treatment of conditions which
 require the inhibition of an enzyme whose preferred mode of action is to catalyse the
 hydrolysis of an ester functionality;

wherein in formula (I):

A is a 6 membered aromatic or hetero-aromatic ring; and

R¹ is a branched or unbranched alkyl (optionally interrupted by one or more oxygen
 atoms), alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, reduced arylalkyl,
 arylalkenyl, heteroaryl, heteroarylalkyl, heteroarylalkenyl, reduced aryl, reduced
 heteroaryl, reduced heteroarylalkyl or a substituted derivative thereof wherein the
 substituents are one or more independently chosen from the group consisting of
 halogen, alkyl, halosubstituted alkyl, aryl, arylalkyl, heteroaryl, reduced heteroaryl,
 reduced heteroarylalkyl, arylalkoxy, cyano, nitro, -C(O)R⁴, -CO₂R⁴, -SOR⁴, -SO₂R⁴, -
 NR⁶R⁷, -OR⁶, -SR⁶, -C(O)CX¹X²NR⁶R⁷, -C(O)NR⁴R⁵, -C(O)N(OR⁵)R⁶, -NR⁶C(O)R⁴,

-CR⁶(NH₂)CO₂R⁶, -NHCX¹X²CO₂R⁶, -N(OH)C(O)NR⁶R⁷, -N(OH)C(O)R⁴, -NHC(O)NR⁶R⁷, -C(O)NHN⁶R⁷, -C(O)N(OR⁵)R⁶, or a lipid or steroid (natural or synthetic), with the proviso that any hetero atom substituent in R¹ and/or R² must be separated from the exocyclic nitrogen atom by at least two carbon atoms (preferably saturated); and

R² is hydrogen or is a group as defined above for R¹;

and where:-


R⁴ is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, reduced heteroaryl, reduced heteroarylalkyl, -OR⁶, -NHCX¹X²CO₂R⁶ or -NR⁶R⁷;

R⁵ is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, reduced heteroaryl or reduced heteroarylalkyl;

R⁶ and R⁷ are independently selected from hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heteroaryl, reduced heteroaryl, heteroarylalkyl, reduced heteroarylalkyl or -(CH₂)_n(OR⁵)_m wherein n is 1 to 12, and m is 1-3; and

X¹ and X² are independently hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, reduced heteroaryl or reduced heteroarylalkyl.

2. Use according to claim 1 wherein the compound (I) is a compound of formula (II):



(II)

or a pharmaceutically acceptable salt, ester, amide or prodrug thereof,
 5 wherein:-

R¹, R², R⁴, R⁵, R⁶, R⁷, X¹ and X² are as defined above for formula (I);

and

10 R⁸, R⁹, R¹⁰, R¹¹ are each independently hydrogen, halo, hydroxy, amino, nitro, cyano,

or a group R¹, as defined above;

15 or a group R¹²Q where Q is O, CO, CONH, NHCO, S, SO, SO₂, or SO₂NH₂ and R¹² is hydrogen or a group R¹ as defined above;

20 or a group R¹R²N where R¹ and R² are as defined above, with the proviso that any hetero atom substituent in R¹ and/or R² must be separated from the aromatic hetero atom substituent by at least two carbon atoms (preferably saturated).

3. Use according to claim 2 wherein in the compound of formula (II) R¹ is aryl or an aryl alkyl group wherein the alkyl moiety has up to 25 carbon atoms, or an aryl aryl

group; wherein the aryl alkyl group or the aryl aryl group may be separated by a spacer where the spacer can be an ester, amide, O, CH₂, or a ketone;

R² is hydrogen or is a group as defined above for R¹;

R⁸ is hydrogen or fluorine;

R⁹ is lower branched or unbranched alkyl having 1 to 10 carbon atoms; cyclic alkyl having 3 to 10 carbon atoms; haloalkyl; or a halogen;

R¹⁰ is lower branched or unbranched alkyl having 1 to 10 carbon atoms; cyclic alkyl having 3 to 10 carbon atoms; haloalkyl; or a halogen;

R¹¹ is hydrogen, lower branched or unbranched alkyl having 1 to 10 carbon atoms, or halogen.

4. Use according to any of claims 1 to 3 wherein R¹ represents phenyl substituted by a group selected from OR¹³, -COR¹³, CO₂R¹³, SOR¹³, SO₂R¹³, CONR¹³R¹⁴, NR¹⁴C(O)NR¹³, C₁₋₁₀alkyl, C₁₋₁₀alkoxy, haloC₁₋₁₀alkyl, aryl, aryl C₁₋₁₀alkyl, heteroaryl or heteroaryl C₁₋₁₀alkyl; wherein R¹³ and R¹⁴ each independently represents hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, C₃₋₆cycloalkyl, C₃₋₆cycloalkenyl, aryl, arylC₁₋₁₀alkyl, heteroaryl, heteroarylC₁₋₁₀alkyl, reduced heteroaryl or reduced heteroarylC₁₋₁₀alkyl.

5. Use according to any of claims 1 to 4 wherein R¹ represents phenyl substituted by OR¹³ or COR¹³ wherein R¹³ is phenyl; phenyl substituted by -CO₂R¹³ wherein R¹³ represents C₁₋₁₀alkyl; or phenyl substituted by C₆₋₁₀alkyl.

6. Use according to any of claims 1 to 5 wherein the enzyme is a lipase.

7. Use according to any of claims 1 to 6 wherein said condition is selected from obesity, hyperlipaemia, hyperlipidaemia, hyperglycaemia (type II diabetes), hypertension, cardiovascular disease, stroke, gastrointestinal disease and gastrointestinal conditions.

5

8. Use according to any of claims 1 to 6 wherein said medicament is for reducing levels of toxins in body fat.


9. Use according to any of claims 1 to 8 wherein the medicament is for administration to humans.

10

10. Use according to any of claims 1 to 8 wherein the medicament is for administration to animals.

11. A compound of formula (IIa):

15



20

(IIa)

or a pharmaceutically acceptable salt, ester, amide or prodrug thereof;

wherein:-

25

R^{1a} represents

(i) a C₁₀₋₃₀ branched or unbranched alkyl, C₂₋₃₀ alkenyl, C₂₋₃₀ alkynyl, cycloalkenyl, aryl-C₁₀₋₃₀ alkyl, aryl-C₁₀₋₃₀ alkenyl, heteroaryl, heteroaryl-C₁₋₃₀ alkyl, heteroaryl-C₂₋₃₀

alkenyl, reduced aryl, reduced heteroaryl, reduced heteroaryl-C₁₋₃₀ alkyl or a substituted derivative thereof wherein the substituents are one or more independently chosen from the group consisting of halogen, C₁₋₁₀ alkyl, halosubstituted C₁₋₁₀ alkyl, aryl, aryl-C₁₋₁₀ alkyl, heteroaryl, reduced heteroaryl, reduced heteroaryl-C₁₋₁₀ alkyl, aryl-C₁₋₁₀ alkoxy, cyano, nitro, -C(O)R¹³, -CO₂R¹³, -SOR¹³, -SO₂R¹³, -NR¹³R¹⁴, -OR¹³, -SR¹³, -C(O)NR¹³R¹⁴, and -NR¹⁴C(O)R¹³, with the proviso that any hetero atom substituent in R¹ must be separated from the exocyclic nitrogen atom by at least two carbon atoms (preferably saturated); or

(ii) aryl substituted by one or more independently chosen from the group consisting of halosubstituted C₁₋₁₀ alkyl, aryl, aryl-C₁₋₁₀ alkyl, heteroaryl, reduced heteroaryl, reduced heteroaryl-C₁₋₁₀ alkyl, aryl-C₁₋₁₀ alkoxy, cyano, -C(O)R¹³, -CO₂R¹³, -SOR¹³, -SO₂R¹³, -NR¹³R¹⁴, -OR¹³ (providing that in this instance R¹³ does not represent aryl or alkyl), -SR¹³, -C(O)NR¹³R¹⁴, and -NR¹⁴C(O)R¹³

wherein:-

R¹³ and R¹⁴ each independently represents hydrogen, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, C₃₋₆cycloalkyl, C₃₋₆cycloalkenyl, aryl, arylC₁₋₁₀alkyl, heteroaryl, heteroarylC₁₋₁₀alkyl, reduced heteroaryl or reduced heteroarylC₁₋₁₀alkyl;

R^{2a} is hydrogen or is a group as defined above for R¹; and

R^{8a}, R^{9a}, R^{10a}, R^{11a} are as defined above for R⁸, R⁹, R¹⁰ and R¹¹ in formula (II).

provided that:

when R^{1a} represents a heteroaryl group it is not thiadiazolyl, triazolyl or thiazolyl; and
when R^{1a} represents a reduced heteroaryl group it is not thiazolidinyl.

12. A compound according to claim 11 wherein R^{1a} represents phenyl substituted by a group selected from OR^{13} (providing in this instance R^{13} does not represent alkyl or aryl), $-COR^{13}$, CO_2R^{13} , SOR^{13} , SO_2R^{13} , $CONR^{13}R^{14}$, $NR^{14}C(O)NR^{13}$, $haloC_{1-10}alkyl$, aryl, aryl $C_{1-10}alkyl$, heteroaryl or heteroaryl $C_{1-10}alkyl$.


5

13. A compound according to claim 11 or claim 12 wherein R^{2a} represents hydrogen or $C_{1-10}alkyl$.

14. A compound according to any of claims 11 to 13 wherein R^{8a} , R^{9a} , R^{10a} and R^{11a} each independently represents hydrogen, halo, hydroxy, amino, nitro, cyano, thiol, $C_{1-10}alkyl$, $C_{1-10}alkoxy$, $C_{1-10}cycloalkyl$, $C_{1-10}cycloalkoxy$, $C(O)R^{15}$, $C(O)NR^{15}R^{16}$, $S(O)R^{4a}$ or $haloC_{1-10}alkyl$; wherein R^{15} and R^{16} each independently represent hydrogen or $C_{1-10}alkyl$.

15

15. A compound of formula (IIb)



20

wherein

R^8 - R^{11} are as defined hereinbefore and

25

R^{20} represents $C_{1-20}alkyl$, $C_{1-20}alkoxy$, or optionally substituted phenoxy.

16. A compound of formula (II) selected from:

30

2-Phenylamino-4*H*-3,1-benzoxazin-4-one;

- 6-Chloro-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
2-Butylamino-4*H*-3,1-benzoxazin-4-one;
6-Methyl-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
2-(4-Chlorophenylamino)-4*H*-3,1-benzoxazin-4-one;
5 6-Methyl-2-(naphth-2-ylamino)-4*H*-3,1-benzoxazin-4-one;
2-Ethylamino-4*H*-3,1-benzoxazin-4-one;
7-Fluoro-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
5-Fluoro-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
7-Methyl-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
10 7-Ethyl-2-Phenylamino-4*H*-3,1-benzoxazin-4-one;
7-Octyl-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
7-Butyl-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
2-Phenylamino-4*H*-pyrido[2,3-*d*][1,3]oxazin-4-one;
6-Nitro-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
15 6-Acetamido-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
2-Phenylamino-7-trifluoromethyl-4*H*-3,1-benzoxazin-4-one;
7-Amino-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
2-Phenylamino-4*H*-pyrido[3,4-*d*][1,3]oxazin-4-one;
2-Cyclopropylamino-4*H*-3,1-benzoxazin-4-one;
20 2-(Naphth-2-ylamino)-4*H*-3,1-benzoxazin-4-one;
2-(6-Phenylhexylamino)-4*H*-3,1-benzoxazin-4-one;
6-Cyano-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
6-Trifluoromethyl-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
6-Formyl-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
25 2-Phenylamino-4*H*-3,1-benzoxazin-4-one-6-sulphinic acid;
7-Hydroxy-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
7-Cyclopropyl-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
6,7-Dimethyl-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
6-Iodo-2-octylamino-4*H*-3,1-benzoxazin-4-one;
30 7-Butyl-2-octylamino-4*H*-3,1-benzoxazin-4-one;
6-Methyl-2-(dodeca-7-ynylamino)-4*H*-3,1-benzoxazin-4-one;
8-Fluoro-2-phenylamino-4*H*-3,1-benzoxazin-4-one;
6-Cyclopropyl-2-phenyl-4*H*-3,1-benzoxazin-4-one;
6-Mercapto-2-phenylamino-4*H*-3,1-benzoxazin-4-one; and
35 6-Cyano-2-phenylamino-4*H*-3,1-benzoxazin-4-one;

or a salt, ester, amide or prodrug thereof.

- 40 17. A compound of formula (IIa) selected from:

2-(4-Trifluoromethylphenylamino)-4*H*-3,1-benzoxazin-4-one;
2-(3-Trifluoromethylphenylamino)-4*H*-3,1-benzoxazin-4-one;
2-(4-Butoxycarbonylphenylamino)-6-methyl-4*H*-3,1-benzoxazin-4-one;

- 2-Hexadecylamino-6-methyl-4*H*-3,1-benzoxazin-4-one;
2-(4-Benzoylphenylamino)-7-methyl-4*H*-3,1-benzoxazin-4-one;
2-(4-Phenoxyphenylamino)-7-trifluoromethyl-4*H*-3,1-benzoxazin-4-one;
2-(2-Cyanophenylamino)-7-methyl-4*H*-3,1-benzoxazin-4-one;
5 2-(3-Cyanophenylamino)-7-methyl-4*H*-3,1-benzoxazin-4-one;
2-(4-Cyanophenylamino)-7-methyl-4*H*-3,1-benzoxazin-4-one;
2-(4-Cyanophenylamino)-4*H*-3,1-benzoxazin-4-one;
2-(4-Carboxyphenylamino)-4*H*-3,1-benzoxazin-4-one;
2-(4-Aminophenylamino)-4*H*-3,1-benzoxazin-4-one;
10 2-(4-Hydroxyphenylamino)-4*H*-3,1-benzoxazin-4-one;
2-(4-*N*-Methylcarbamoylphenylamino)-4*H*-3,1-benzoxazin-4-one;
2,2'-(1,8-Octyldenediamino)-*bis*-4*H*-3,1-benzoxazin-4-one;
2-(Pyrrol-3-ylamino)-4*H*-3,1-benzoxazin-4-one;
2-(Piperidin-4-ylamino)-4*H*-3,1-benzoxazin-4-one;
15 2-[6-(Pyrrol-2-yl)-hexylamino]-4*H*-3,1-benzoxazin-4-one;
2-(4-Ethoxycarbonylphenylamino)-4*H*-3,1-benzoxazin-4-one; and
6-Methyl-2-[6-(thien-2-yl)hexylamino]-4*H*-3,1-benzoxazin-4-one;

or a salt, ester, amide or prodrug thereof.

20

18. A compound of formula (IIb) selected from:


- 2-(4-Butylphenylamino)-4*H*-3,1-benzoxazin-4-one;
2-(4-Methoxyphenylamino)-4*H*-3,1-benzoxazin-4-one;
2-(4-Methylphenylamino)-4*H*-3,1-benzoxazin-4-one;
25 2-(4-Phenoxyphenylamino)-4*H*-3,1-benzoxazin-4-one;
2-[4(1-Methylethyl)phenylamino]-4*H*-3,1-benzoxazin-4-one;
6-Methyl-2-(4-phenoxyphenylamino)-4*H*-3,1-benzoxazin-4-one;
7-Ethyl-2-Phenylamino-4*H*-3,1-benzoxazin-4-one;
2-(4-Hexylphenylamino)-6-methyl-4*H*-3,1-benzoxazin-4-one;
30 2-(4-Heptyloxyphenylamino)-6-methyl-4*H*-3,1-benzoxazin-4-one;
7-Methyl-2-(4-phenoxyphenylamino)-4*H*-3,1-benzoxazin-4-one;
7-Methyl-2-(2-phenoxyphenylamino)-4*H*-3,1-benzoxazin-4-one;
7-Methyl-2-(3-phenoxyphenylamino)-4*H*-3,1-benzoxazin-4-one;
7-Methyl-2-(4-octylphenylamino)-4*H*-3,1-benzoxazin-4-one;
35 2-(2-Phenoxyphenylamino)-4*H*-3,1-benzoxazin-4-one; and
2-(3-Phenoxyphenylamino)-4*H*-3,1-benzoxazin-4-one;

or a salt, ester, amide or prodrug thereof.

40

19. A process for the preparation of a novel compound of formula (II) in particular a compound of formula (IIa) or formula (IIb) which process comprises:


Process (A) cyclising a compound of formula (III)



10 wherein R^1 and R^8 - R^{11} are as hereinbefore defined and R^{18} is hydrogen or C_{1-6} alkyl.

or:

15 Process (B) reacting a compound of formula (IV)



with an amine of formula (V)



or:

30 Process (C) converting a compound of formula (I), (II), (IIa) or (IIb) into a different compound of formula (IIa) or (IIb), by, for example,

(i) reduction of a compound of formula (I), (II), (IIa) or (IIb) wherein any of R^1 , R^8 , R^9 , R^{10} and R^{11} contains an alkenyl or alkynyl group or moiety, to the corresponding alkyl or alkenyl group or moiety; or

5 (ii) alkylation of a compound of formula (I), (II), (IIa) or (IIb) where one or more of R^8 , R^9 , R^{10} and R^{11} represents a halogen atom.

20. A compound, as claimed in any one of claims 11 to 18, for use in medicine.

10 21. A pharmaceutical composition comprising a novel compound of formula (II) or a pharmaceutically acceptable salt, ester, amide or pro-drug thereof, in combination with a pharmaceutically acceptable carrier or diluent.

15 22. A food product comprising a compound of formula (II) or a pharmaceutically acceptable salt, ester, amide or pro-drug thereof.

23. A method for the prevention or treatment of obesity or an obesity related disorder, the method comprising administering a compound, as defined in any one of claims 1 to 18, or a composition as claimed in claim 21 or claim 22 to a patient.

20 24. A compound, as defined in any one of claims 1 to 18, for use in the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality.

25 25. Use of a compound as defined in any of claims 1 to 18 or a pharmaceutically acceptable salt, ester, amide or prodrug thereof to reduce fat content of animals which provide meat for human consumption.

26. A cosmetic method for maintaining a given weight, or for cosmetic weight loss, the method comprising the administration of a compound as defined in any of claims 1 to 18.

5 27. A compound comprising formula (I) hereinbefore described with reference to one or more of the examples.

28. A process for obtaining a compound comprising formula (I) hereinbefore described with reference to one or more of the examples.

10

29. The use of a compound comprising formula (I) in the inhibition of an enzyme whose preferred mode of action is to catalyse the hydrolysis of an ester functionality hereinbefore described with reference to one or more of the examples.

15 30. Use according to claim 29 of a compound of formula (I) in the control and inhibition of unwated enzymes in a process or product.

31. Use according to claim 30 in the manufacture of healthcare goods comprising surfactants, soap or detergents.

20

32. Use according to claim 30 in preventing the degradation of foodstuff which comprises a fat.

25

30

INTERNATIONAL SEARCH REPORT

I. International Application No
PCT/GB 00/00032

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/536 A61P3/04 C07D265/24 C07D498/04 C07D413/12
/(C07D498/04, 265:00, 221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	M. GÜTSCHOW ET AL.: "Inhibition of cathepsin G by 4H-3,1-benzoxazin-4-ones" BIOORGANIC & MEDICINAL CHEMISTRY, vol. 5, no. 10, 1997, pages 1935-1942, XP000904822 the whole document	1-27
X	R.L. JARVEST ET AL.: "Inhibition of HSV-1 protease by benzoxazinones" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, vol. 6, no. 20, 1996, pages 2463-2466, XP000904816 the whole document	1-27
X	WO 96 37485 A (SEARLE & CO ;ABOOD NORMAN (US); FLYNN DANIEL L (US); BECKER DANIEL) 28 November 1996 (1996-11-28) claims	1-27
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

18 April 2000

Date of mailing of the international search report

28/04/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Chouly, J

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/00032

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 07648 A (WARNER LAMBERT CO) 14 March 1996 (1996-03-14) claims & US 5 652 237 A cited in the application ---	1-27
X	EP 0 147 211 A (SYNTEX INC) 3 July 1985 (1985-07-03) claims & US 4 657 893 A cited in the application ---	1-27
X	DE 23 15 303 A (BAYER AG) 17 October 1974 (1974-10-17) cited in the application claims ---	1,19
X	US 3 450 700 A (ULRICH HENRI ET AL) 17 June 1969 (1969-06-17) the whole document -----	1-27

INTERNATIONAL SEARCH REPORT

international application No.

PCT/GB 00/ 00032

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 23,26
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 23,26
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☒ Claims Nos.: 27-32
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
Claims 27-32 were not searched since they refer to any compounds,
formulations, processes and methods mentioned in the description;
therefore it does not comply with the requirements of the PCT, Rule 6.2a.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/GB 00 00032

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 27-32

Claims 27-32 were not searched since they refer to any compounds, formulations, processes and methods mentioned in the description; therefore it does not comply with the requirements of the PCT, Rule 6.2a.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/00032

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9637485	A	28-11-1996	US 5985872 A	16-11-1999
			AU 711174 B	07-10-1999
			AU 5803096 A	11-12-1996
			BR 9608506 A	15-06-1999
			CA 2222034 A	28-11-1996
			CN 1190963 A	19-08-1998
			EP 0828721 A	18-03-1998
			JP 11505849 T	25-05-1999
			NZ 308398 A	29-06-1999
WO 9607648	A	14-03-1996	US 5652237 A	29-07-1997
			AU 3496295 A	27-03-1996
EP 0147211	A	03-07-1985	US 4657893 A	14-04-1987
			AT 56444 T	15-09-1990
			AU 586616 B	20-07-1989
			AU 3716984 A	04-07-1985
			CA 1269800 A	29-05-1990
			DK 625184 A	28-06-1985
			ES 539038 D	01-06-1986
			ES 8607260 A	01-11-1986
			ES 550879 D	01-03-1987
			ES 8703437 A	01-05-1987
			FI 845116 A,B,	28-06-1985
			HU 36808 A	28-10-1985
			IL 73943 A	31-01-1989
			JP 60169469 A	02-09-1985
			KR 8900992 B	15-04-1989
			NO 845176 A,C	28-06-1985
			NZ 210669 A	30-05-1988
			ZA 8410089 A	27-08-1986
DE 2315303	A	17-10-1974	NONE	
US 3450700	A	17-06-1969	NONE	